

REVIEW

Collective Cell Migration of the Cephalic Neural Crest: The Art of Integrating Information

Eric Theveneau and Roberto Mayor*

Department of Cell and Developmental Biology, University College London, United Kingdom

Received 5 November 2010; Revised 30 November 2010; Accepted 4 December 2010

Summary: The cephalic neural crest (NC) cells delaminate from the neuroepithelium in large numbers and undergo collective cell migration under the influence of multiple factors including positive and negative taxis, cell–cell interactions mediating cell sorting, cell cooperation, and Contact-Inhibition of Locomotion. The migration has to be tightly regulated to allow NC cells to reach precise locations in order to contribute to various craniofacial structures such as the skeletal and peripheral nervous systems. Several birth defects, syndromes, and malformations are due to improper cephalic NC (CNC) migration, and NC cell migration bears important similarities to cancer cell invasion and metastasis dissemination. Therefore, understanding how CNC cells interpret multiple inputs to achieve directional collective cell migration will shed light on pathological situations where cell migration is involved. *genesis* 49:164–176, 2011. © 2010 Wiley-Liss, Inc.

Key words: neural crest cells; collective cell migration; chemotaxis; Contact-Inhibition of Locomotion; craniofacial structures

INTRODUCTION

The neural crest (NC) is a multipotent cell population induced at the neural plate border and located later in the dorsal part of the neural tube. Once specified, the NC cells delaminate from the neuroepithelium and migrate throughout the embryo following well-defined routes to their final locations where they stop and differentiate in a wide range of cell types (Hall, 2008; Le Douarin and Kalcheim, 1999). This review will focus on the cephalic NC (CNC) population, which arises from the diencephalon to the caudal hindbrain. The CNC makes an outstanding contribution to the craniofacial structures, and several syndromes and birth defects are due to improper CNC migration. Understanding how

CNC migration is precisely regulated in time and space gives precious information about NC-related diseases and malformations. In addition, it sheds light on the general mechanisms involved in regulating cell migration in normal and pathological situations. Here, we present a brief overview of the CNC contribution to the craniofacial structures and the main signals regulating CNC migration and discuss how cells can integrate multiple and somehow opposite inputs to achieve directional migration.

CONTRIBUTION OF THE CNC TO THE CRANIOFACIAL STRUCTURES

The CNC makes critical contributions to the cranial nerves (CN) and ganglia, the skeletal system (including teeth and ossicles of the middle ear), the ocular and perocular structures, smooth muscles, and connective tissues of the blood vessels. In addition, it forms the dermis of the head, most of the pigment cells (iris cells are not NC derivatives), and the meninges of the proencephalon as well as influencing striated muscle patterning and the migration of placode-derived neurones (Creuzet *et al.*, 2005; Hall, 2008; Johnston *et al.*, 1979; Le Douarin and Kalcheim, 1999; Noden and Trainor, 2005).

Cephalic Peripheral Nervous System

The CNC produces the glial cells (Schwann cells) of all CN and ganglia except for the CN II, which is ensheathed in the meningeal layers, myelinated by oligodendrocytes, and lies in the central nervous system

* Correspondence to: Roberto Mayor, Department of Cell and Developmental Biology, University College London, United Kingdom.

E-mail: r.mayor@ucl.ac.uk

Contract grant sponsor: MRC, BBSRC, and Wellcome Trust

Published online 14 December 2010 in

Wiley Online Library (wileyonlinelibrary.com).

DOI: 10.1002/dvg.20700

(Schoenwolf *et al.*, 2009). In addition, CNC cells give rise to neurons of the proximal ganglia of CN VII (facial), IX (glossopharyngeal), and X (vagus). Alongside with ectodermal placodes, they also contribute to neuroones of the ciliary (CN III/oculomotor), trigeminal (CN V/trigeminal), and vestibular (CN VIII/vestibulocochlear) ganglia. Distal ganglia of the CN VII (geniculate), IX (petrosal), and X (nodose) as well as the acoustic ganglion contain neurones of placodal origin (summarized in Fig. 1A–C). Therefore, neuronal CNC derivatives are not only involved in controlling eye movements (CN III), biting, chewing, and swallowing (CN V) and facial expressions (CN VII) but also carried sensory information from, for instance, the face, nasal cavity and mouth (CN V), the sense of taste (CN VII and IX), and the sense of balance (CN VIII). Finally, they relay information about the state of the body's organs to the central nervous system through the vagus nerve (CN X) (D'Amico-Martel and Noden, 1983; Hall, 2008; Le Douarin and Kalcheim, 1999; Lee *et al.*, 2003; Schoenwolf *et al.*, 2009).

Skeletal System: Skull, Tooth, and Ear Ossicles

Studies using amphibians, quail-chick chimeras, and transgenic mice have highlighted the fact that CNC cells contribute to most of the cartilages and bones of the head and neck (Fig. 1D) (Couly *et al.*, 1993; Evans and Noden, 2006; Gross and Hanken, 2005; Hall, 2008; Hanken and Gross, 2005; Helms and Schneider, 2003; Helms *et al.*, 2005; Le Douarin and Kalcheim, 1999; Minoux and Rijli, 2010; Morriss-Kay, 2001; Trainor, 2005; Trainor and Tam, 1995). Although it is well accepted that all facial skeletal structures are NC-derived, the limit between NC-derived and mesoderm-derived skeletons at the back of the skull and the jaw are still matters of debate. Although studies on avian models classify the parietal bone as a NC derivative, in mammals, it is considered as mesoderm-derived. In addition, mice data support the idea that part of the jaw may be formed by non-NC cells, whereas avian data find it to be entirely NC-derived. A possible explanation for these discrepancies may be that different technical approaches were used. In mice, CNC derivatives were mapped using a mouse model expressing the gene-encoding β -galactosidase under the control of Wnt1 promoter, leaving the possibility that some NC subpopulations may not express Wnt1. In birds, however, the quail-chick chimera system provides a stable means of staining cells and labels all derivatives regardless of the fluctuations of gene expressions among subpopulations. An alternative explanation would be that in mammals some of the NC prerogatives have been transferred to the mesoderm. For a precise account of studies about CNC contribution to the skull in different animal models and discussion about the origin of the parietal bone

and the jaw, the reader will find useful information in the references cited earlier. Apart from its contribution to the skull, the CNC gives rise to ossicles of the middle ear (Gross and Hanken, 2008; Hall, 2008; Le Douarin and Kalcheim, 1999; Le Lievre, 1978) (Fig. 1E), which transmit vibrations from the tympanic membrane to the cochlea. Finally, NC cells also form multiple parts of the teeth (Fig. 1F) including the dentin that is deposited and mineralized by odontoblasts of NC origin (Graveson *et al.*, 1997; Lumsden, 1988; Lumsden and Buchanan, 1986; Smith and Hall, 1990).

Neurocristopathies

Syndromes, tumors, and malformations due to incorrect NC development are named neurocristopathies. Because of their implications in a wide variety of craniofacial structures, problems occurring during CNC formation, migration, and differentiation have dramatic consequences on the development of the head. Induction, proliferation, or survival issues usually give rise to dysplasia (abnormal development of an organ or part of the body, including congenital absence) such as seen in Treacher Collins syndrome (Trainor, 2010; Walker and Trainor, 2006), while CNC migration defects lead to malformations. Cleft lip, cleft palate, defects of the anterior chamber of the eye, unusual (or lack of) pigmentation, and abnormal ear development are common features of CNC migration defects that can be found in diseases like the CHARGE association and Waardenburg, DiGeorge, and Goldenhar syndromes (Cohen, 1989, 1990; Hall, 2008; Jones, 1990; Le Douarin and Kalcheim, 1999; Schoenwolf *et al.*, 2009; Wurdak *et al.*, 2006).

CNC CELL MIGRATION

All along the antero-posterior axis NC cells first separate themselves from the neuroepithelium during the delamination phase and then start migrating following stereotypical routes. Although the general sequence of induction, delamination, and migration is true for all NC cells emerging along the antero-posterior axis, CNC cells bear some specific features that are worth noting. Although trunk NC cells delaminate progressively in a dripping fashion over a long period, CNC cells depart almost all at once (Hall, 2008; Le Douarin and Kalcheim, 1999; Theveneau *et al.*, 2007). Therefore, they start migrating as a multilayered cell group, whereas trunk NC cells exhibit a single chainlike organization. The delamination at the cephalic level is regulated by numerous transcription factors including *snail2*, *lsox5*, and *ets1*, but our understanding of their relationships remains incomplete (del Barrio and Nieto, 2002; Perez-Alcala *et al.*, 2004; Taneyhill *et al.*, 2007; Theveneau *et al.*, 2007). This process involves a change of cadherin expression reducing cell-cell adhesion, which, in chick,




FIG. 1. Neural crest contribution to the craniofacial structures. **A:** Neural crest cells neuronal contribution to cranial PNS shown in a virtual mammalian embryo. **B:** Distribution of the cranial nerves in a human head. Nerves receiving neuronal contribution from the neural crest are in color. The nerves receiving only glial contribution, or no NC contribution (CN II, see text), appear in gray. **C:** Detailed neuronal contribution of neural crest and placodal cells in cranial ganglia III, V, VII, VIII, IX, and X. Origin of the CNC contribution along the antero-posterior axis is indicated under each ganglion. **D:** Human head showing the global repartition of neural crest and mesoderm-derived bones and cartilages. Only the main NC-derived facial elements are labeled. **E:** Neural crest cells form the three ossicles of the middle ear. **F:** Contribution of the neural crest to various parts of the teeth. mes, mesencephalon; r, rhombomere.

involves a global shift from type I (N-Cadherin) to type II (Cadherin 6B, 7) cadherin. However, although migrating NC express reduced levels of type I cadherin, these molecules are still present as shown by the expression

of N-cadherin in *Xenopus*, zebrafish, and chick (Piloto and Schilling, 2010; Theveneau *et al.*, 2007, 2010). In addition, CNC cells migrate almost exclusively in close association with the surface ectoderm, at least for the




FIG. 2. Molecular control of neural crest migration. **A:** Overview of the cephalic neural crest migration. CNC cells invade their surrounding tissues in register with their original location along the AP axis. **B:** Detailed view of the early phase of CNC migration during which anterior and posterior migration of cells from r3 and r5 give rise to NC-free regions adjacent to the neural tube. **C:** A stream of NC cells migrating through the extracellular matrix. Cells are polarized according to their cell contact. Inhibitors keeping streams separated are shown in red, and attractant or permissive molecules are shown in green. Numbers refer to panels shown hereafter. Overview of the molecules involved in cell-matrix interactions (1), chemotaxis and chemokinesis (2), defining NC routes borders (3), contact-inhibition of locomotion (4), and cell sorting (5) during CNC cells migration. ant, anterior; di, diencephalon; FGF, fibroblast growth factor; Fn, fibronectin; mes, mesencephalon; Npn1, neuropilin-1; PDGF, platelet-derived growth factor; post, posterior; r, rhombomere; Sdf1, stromal cell-derived factor-1; Syn4, syndecan-4; VEGF, vascular endothelium growth factor.

first part of their migration, whereas trunk NC cells split into several subpopulations migrating deep inside the embryo toward the dorsal aorta through the somites and finally underneath the ectoderm (Kuriyama and

Mayor, 2008; Le Douarin and Kalcheim, 1999). Guidance of collective CNC cell migration involves various molecules, and signaling pathways are summarized in Figure 2 and discussed below.

From a Continuum to Discrete Streams

Despite delaminating continuously from the diencephalon to the caudal hindbrain, the CNC will quickly split into the three following streams (Fig. 2A,B): adjacent to the neural tube from the diencephalon to r2, adjacent to r4, and adjacent to the postotic region from r6 to r8. Crests from r3 and r5 migrate rostrally and caudally to join adjacent streams leaving two NC-free zones opposite to r3 and r5 (Kulesa *et al.*, 2010; Kuriyama and Mayor, 2008; Lumsden and Guthrie, 1991; Lumsden *et al.*, 1991; Sechrist *et al.*, 1993). Several signaling pathways control the separation of the original NC continuum into discrete groups (Figs. 2B,C.3). First of all, molecules of the class 3-semaphorins family (3A, 3F and 3G) and their neuropilin receptors (Npn1 and 2) are involved in keeping NC-free zones adjacent to r3 and r5 in chick, mouse, and zebrafish (Eickholt *et al.*, 1999; Gammill *et al.*, 2007; Osborne *et al.*, 2005; Schwarz *et al.*, 2008; Yu and Moens, 2005). CNC cells express the neuropilin receptors while surrounding tissues express and secrete soluble semaphorin ligands (Fig. 2C, red). Inhibition of the semaphorin signaling causes ectopic NC cells to invade spaces in between the normal streams. In addition, *in vivo* experiments and *in vitro* assays on stripes of semaphorin ligands showed that NC cells preferentially avoid semaphorin-containing regions. In *Xenopus*, CNC cells also express Neuropilins during migration (Koestner *et al.*, 2008), but the role of Semaphorin signaling remains unexplored.

Besides Semaphorin signaling, transmembrane ephrins and Eph receptors also play a role in this process (Figs. 2C.3,C.5). Studies in *Xenopus*, mice, and chick have highlighted a great variability in the combination of the ephrins and Eph expressed by CNC and the head mesoderm, but their functions seem conserved (Adams *et al.*, 2001; Davy *et al.*, 2004; Mellott and Burke, 2008; Smith *et al.*, 1997). Loss- and gain-of-function experiments lead to two main phenotypes: invasion of NC-free zones or misaddressing of NC cells into an unusual stream without abolishing NC-free zones. Therefore, ephrins and Eph seem to play a dual role. They are involved in cell sorting, helping place CNC into a specific stream according to their antero-posterior level of origin, and they also prevent entry into specific areas (Fig. 2C.3,C.5).

Finally, Slit ligands and their Robo receptors (Dickson and Gilestro, 2006; Ypsilanti *et al.*, 2010) have been implicated in regulating trunk NC-cell migration (De Bellard *et al.*, 2003; Jia *et al.*, 2005; Kuriyama and Mayor, 2008) by driving NC cells into the ventromedial pathway. However, at cephalic levels, while Slit/Robo signaling seems to control late events related to ganglia formation (Shiau and Bronner-Fraser, 2009), its putative role in guidance of CNC migration has yet to be assessed.

Chemotaxis, Chemokinesis: New Findings for Old Ideas

Until very recently, the accepted model for NC cell migration was based on the fact that NC cells were mesenchymal and highly migratory cells facing permissive areas containing extracellular matrix (ECM) and restrictive areas defined by negative cues present in the local environment. Consequently, NC cells would explore their direct surroundings and invade all areas free of inhibitors. Positive taxis attracting NC to specific locations and negative taxis repelling NC from the neuroepithelium were proposed (Erickson, 1985, 1988; Erickson and Olivier, 1983; Sechrist *et al.*, 1994) but remained unsupported or, in the case of negative taxis, were ruled out by experimental data. However, over the last few years, several molecules have been shown as targeting trunk NC cells to specific areas (Druckendro and Epstein, 2007; Jiang *et al.*, 2003; Santiago and Erickson, 2002) and more recently attracting or positively influencing CNC migration (Fig. 2C.2). Stromal-cell-derived factor 1 (Sdf1/Cxcl12), a potent chemoattractant for germ cells and lymphocytes, among other cell types (Burger and Kipps, 2006; David *et al.*, 2002; Ganju *et al.*, 1998; Molyneaux *et al.*, 2003; Raz and Mahabaleshwar, 2009; Stebler *et al.*, 2004), was described as an attractant for trunk NC cells invading the hair follicles and required for the formation of the dorsal root ganglia (Belmadani *et al.*, 2005, 2009). Moreover, it has been shown as being required for CNC migration in zebrafish (Olesnick Killian *et al.*, 2009). In addition, in *Xenopus laevis*, we demonstrated that Sdf1 acts as a strong chemoattractant *in vitro* and that its expression is required for normal migration *in vivo*. Moreover, Sdf1 misexpression leads to ectopic CNC invasion in between CNC streams or induces an early arrest of migration (Theveneau *et al.*, 2010). Sdf1/Cxcr4 axis functions primarily by increasing Rac1 activity, a small Rho GTPase required for lamellipodia formation, and stabilizing cell protrusions, which generate directional movement toward the Sdf1 source. Finally, in chick, expression of Sdf1 and its receptor Cxcr4 suggests that Sdf1 may play a similar role in the CNC, but this remains to be addressed (Rehimi *et al.*, 2008; Yusuf *et al.*, 2005).

Besides Sdf1/Cxcr4 axis, VEGF, PDGF, and FGF-signaling pathways have also been involved in positively regulating CNC migration (Fig. 2C.2). CNC cells in chick, *Xenopus*, zebrafish, and mouse express PDGF receptor α (Ho *et al.*, 1994; Le Douarin and Kalcheim, 1999; Orr-Urtreger and Lonai, 1992; Orr-Urtreger *et al.*, 1992; Schatteman *et al.*, 1992; Takakura *et al.*, 1997). In zebrafish, PDGF signaling attracts some CNC into the oral region while in mice it is believed to be involved in conferring migratory abilities on the cells. Data on PDGF signaling during CNC migration are scarce, and its precise influence on cell motility or directionality remains elusive (for a recent and complete summary of PDGF roles in NC development, see Smith and Tallquist, 2010).

VEGF and FGF signaling are important for CNC homing into the second branchial arch in chick and mouse, respectively (McLennan *et al.*, 2010; Trokovic *et al.*, 2005). However, whether or not they act as CNC chemoattractants is unclear. Ectopic VEGF leads to a slight change of CNC migration but fails to induce ectopic NC migration, and FGF-based chemoattraction has not been directly tested yet. Furthermore, the homogenous distribution of VEGF and FGF ligands along the pathways of migration makes it difficult to postulate that they give clear directional information. Alternatively, they may promote general random motility (chemokinesis) rather than chemotaxis.

Cell-Cell Interactions: A Touching Story

CNC cells delaminate in large numbers (Hall, 2008; Le Douarin and Kalcheim, 1999; Theveneau *et al.*, 2007) and therefore encounter a very high cell density at the beginning of their migration. Despite this situation, the fact that CNC cells may influence each other's behavior because of direct interactions has been globally overlooked, even though the idea of Contact-Inhibition of Locomotion, a process during which two cells collapse their protrusions when contacting each other (Abercrombie and Heaysman, 1953; Mayor and Carmona-Fontaine, 2010), was proposed as a driving force for the migration of trunk NC cells decades ago (Erickson, 1988). Recent observations and experimentations made in chick, *Xenopus*, and *zebrafish* embryos showed that cell-cell contacts between two migratory NC cells have a direct influence on how cells actually move (Carmona-Fontaine *et al.*, 2008; Kulesa and Fraser, 2000; Teddy and Kulesa, 2004; Theveneau *et al.*, 2010). Original *in vivo* observations in chick by Kulesa and colleagues (Kulesa and Fraser, 2000; Teddy and Kulesa, 2004) showed two types of response after cell-cell interactions. Cells can retract their protrusions upon contact and stop migrating for a while before resuming migration, or, alternatively, a cell can touch another cell located just in front and follow it. In *Xenopus* embryo, we demonstrated that Contact-Inhibition of Locomotion is taking place when two CNC cells are in contact (Carmona-Fontaine *et al.*, 2008). Among large groups of CNC cells, Contact-Inhibition of Locomotion prevents the formation of cell protrusions in between the cells (Fig. 3A,B1). However, at the border of a NC cluster, leading cells are free of Contact-Inhibition of Locomotion, and they can form protrusions away from the cluster (Fig. 3B2). At the cell-cell contact, cell protrusions collapse, while new protrusions are formed at the opposite end of the cell. We showed that Contact-Inhibition of Locomotion is dependant on N-Cadherin-mediated cell-cell interaction (Theveneau *et al.*, 2010), which triggers the noncanonical Wnt/PCP pathway (Figs. 2C.4 and 3D). This leads to RhoA activation (Carmona-Fontaine *et al.*, 2008; Matthews *et al.*, 2008) and

Rac1 inhibition at the cell-cell contact (Theveneau *et al.*, 2010). Interfering with Wnt/PCP or N-Cadherin expression/activity, both suppresses Contact-Inhibition of Locomotion and leads to ectopic protrusions in between the cells and a lack of repolarization upon cell collision (Carmona-Fontaine *et al.*, 2008; Theveneau *et al.*, 2010). We also showed that nonmotile cells (i.e., dividing cells) can be carried passively by their direct motile neighbors (Theveneau *et al.*, 2010). Moreover, cells responding to Sdf1 can help cells that do not express Cxcr4 to reach the source of the chemoattractant (Theveneau *et al.*, 2010). More surprisingly, we found that cell polarity induced by Contact-Inhibition of Locomotion is required for efficient chemotaxis toward Sdf1. Briefly, cell dissociation completely abolished chemotaxis without interfering with cell motility. On the contrary, a progressive increase in cell density can rescue chemotaxis. Altogether, these data indicate that CNC cells undergo collective cell migration and that cell-cell interactions are critical for cell polarity, cell coordination, and chemotaxis during this process (Carmona-Fontaine *et al.*, 2008; Matthews *et al.*, 2008; Mayor and Carmona-Fontaine, 2010; Theveneau *et al.*, 2010).

Metalloproteinases: The NC Footprint

CNC cells receive information from signals released in their local environment and from direct contact with other NC cells or other cell types, but it is likely that they also have a direct impact on the tissues they invade. Studies in chick, *Xenopus*, and mouse showed that CNC cells express various matrix metalloproteinases [MMP2, 8, 14 (MT1-MPP)] (Cai and Brauer, 2002; Cai *et al.*, 2000; Cantemir *et al.*, 2004; Duong and Erickson, 2004; Giambernardi *et al.*, 2001; Harrison *et al.*, 2004; Tomlinson *et al.*, 2009) and ADAM proteins (ADAM 9, 13, and 19) (Alfandari *et al.*, 1997; Cai *et al.*, 1998; Neuner *et al.*, 2009). These enzymes are able to degrade a wide range of molecules and have been implicated in invasive behavior in cancer and normal cell migration (Edwards *et al.*, 2008; Sternlicht and Werb, 2001). From studies done on other systems, MMPs and ADAMs could be involved in many different ways during CNC migration. By degrading ECM components, they could unravel cryptic binding sites for integrins and therefore promote migration (Giannelli *et al.*, 1997; Petitclerc *et al.*, 1999). Such a role in matrix remodeling has been proposed in NC cells to explain why NC migration is affected after MMPs or ADAMs loss-of-function (Alfandari *et al.*, 2001; Duong and Erickson, 2004), but functional experiments to support this idea are still missing *in vivo*. The local degradation of the ECM could, in addition, release growth factors and signaling molecules that are trapped and would be otherwise inaccessible for NC cells. Such a situation has been previously demonstrated for TGF β (Imai *et al.*, 1997). For




FIG. 3. Contact-inhibition and chemotaxis establish and stabilize cell polarity. **A:** Neural crest cells migrating along a gradient of chemoattractant (shades of green). Numbers refer to cells shown in **B**. **B:** Distribution of RhoA (blue) and Rac1 (red) activity in CNC that are surrounded by other cells (inner cells; 1), that have a cell contact and a free edge (outer cell; 2, 3), and CNC that have no contact with other cells (single cells; 4). Cell interactions establish cell polarity along a contact (RhoA)—free edge (Rac1) axis. The chemoattractant reinforces contact-dependent polarity by increasing Rac1 activity at the free edge but is unable to polarize single cells. **C:** Antagonistic relationship between contact-inhibition and chemotaxis signaling inside an outer cell through opposite regulations of small RhoGTPases. **D:** Signalling cascade controlling RhoA and Rac1 activity in *Xenopus* NC cells [after (Carmona-Fontaine *et al.*, 2008; Matthews *et al.*, 2008; Theveneau *et al.*, 2010; Theveneau and Mayor, 2010)].

instance, Sdf1 and members of the FGF, VEGF, and PDGF families interact directly with Fibronectin (Martino and Hubbell, 2010; Pelletier *et al.*, 2000), which can be degraded by MMP2, MT1-MMP, and ADAM13 (Alfandari *et al.*, 2001; Gaultier *et al.*, 2002; Sternlicht and Werb, 2001). Moreover, MMPs and ADAMs could help in shaping gradients out of homogenous expression patterns. Both MMP2 and MT1-MMP can cut and inactivate Sdf1 (McQuibban *et al.*, 2001; Rodriguez *et al.*, 2010). *Xenopus* CNC expresses MT1-MMP (Harrison *et al.*, 2004; Tomlinson *et al.*, 2009). Therefore, cells could digest Sdf1 present in the environment as they migrate, maintaining a sharp gradient at the front of the NC stream and virtually pushing the target as they move forward. Interestingly, some of the signaling pathways involved in CNC cell migration such as Sdf1/CXCR4 or PDGF stimulate MMPs' expression (Busillo and Benovic, 2007; Smith and Tallquist, 2010).

Thus, a complex feedback loop may take place where MMPs and ADAMs could first help NC cells to use the ECM efficiently and access ligands trapped into it and then degrade these molecules to shape gradients. In addition, by degrading molecules, proteinases may eventually get rid of the signals regulating their own expression. Finally, alongside specific changes of cadherin expression mentioned earlier, MMPs and ADAMs may regulate the mesenchymalization and the level of cell-cell interaction of the CNC by degrading Cadherins as it has been proposed for trunk NC cells (Shoval *et al.*, 2007). Indeed, MMP2 and ADAM13 can digest N-Cadherin and Cadherin-11, respectively (Covington *et al.*, 2005, 2006; Hartland *et al.*, 2009; McCusker *et al.*, 2009).

Despite MMPs and ADAMs molecules holding great potential as putative key players in CNC migration, we still have a lot to discover about their precise role. The




FIG. 4. Integration of multiple signals leading to directional collective migration of the neural crest cells. Cell dispersion promoted by contact-inhibition and epithelium-to-mesenchyme transition is restricted by restrictive cues and chemoattractants. These signals ensure high cell motility, high levels of cell interactions, full response to chemoattractants, and altogether drive directional collective cell migration of the cephalic neural crest cells.

recent advances in imaging CNC *in vivo* may help to decipher some of their functions during CNC invasion of the surrounding tissues.

The Art of Integrating Information

CNC may seem overwhelmed by information, part of which could also appear as counter productive. So how does it all make sense?

Previous studies in chick on enteric NC cells invading the gut and CNC entering the branchial arches and recent work in *Xenopus* CNC have shown that regulation of cell density and cell-cell interactions are critical aspects for efficient NC-cell migration (Barlow *et al.*, 2008; Kulesa *et al.*, 2008; Simpson *et al.*, 2007; Theveneau *et al.*, 2010). In *Xenopus* CNC, reducing cell number leads to fewer cell interactions, a poor or randomized cell polarity, inefficient chemotaxis, and reduced cell spreading. On the other hand, maintaining strong cell-cell adhesion, which may seem the best way to maintain a high cell density, also disrupts CNC migration (Theveneau *et al.*, 2010). Therefore, it is clear that, even if a proper epithelium-to-mesenchyme transition is required to confer full migratory potential on the NC cells, cell density has to be maintained by other means. In light of recent advances, we can propose a model integrating these multiple inputs into a general control of CNC collective cell migration (see Fig. 4). In this model, epithelium-to-mesenchyme transition and

Contact-Inhibition of Locomotion promote cell dispersion (green, Fig. 4), while restrictive cues preventing cells from exiting normal routes and chemoattractants leading to cell accumulation in precise locations both concur to maintain high cell density (red, Fig. 4). This high cell density ensures high levels of cell interactions, cell polarization, and a full response to chemoattractants, which altogether drive directional collective cell migration of the CNC cells (see Fig. 4).

However, for cells to be able to integrate all this information, common effectors are needed at the molecular levels. Interestingly, some of these events are indeed mediated by the same molecules. Neuropilin-1, for instance, acts as a co-factor for both permissive (VEGF) and restrictive (class 3-Semaphorin) signaling (McLennan and Kulesa, 2007, 2010; McLennan *et al.*, 2010). In addition, N-Cadherin activity has to be lowered to allow mesenchymalization, but not abolished, as it is required to mediate Contact-Inhibition of Locomotion (Theveneau *et al.*, 2010). More importantly, some of these signals, namely, Contact-Inhibition of Locomotion and chemotaxis, regulate common intracellular effectors in opposite manners (see Fig. 3). Indeed, while N-Cadherin/Contact-Inhibition of Locomotion lead to RhoA activation, Rac1 inhibition, and protrusion collapse (Carmona-Fontaine *et al.*, 2008; Matthews *et al.*, 2008; Theveneau *et al.*, 2010), and Sdf1 increases Rac1 activity and stabilizes lamellipodia (Theveneau *et al.*, 2010). Finally, Syndecan-4, being able to interact

with Fibronectin and Sdf1, (Carey, 1997; Charnaux *et al.*, 2005; Matthews *et al.*, 2008; Pelletier *et al.*, 2000) could have a dual role in CNC migration. Associated with fibronectin, Syndecan-4 leads to Rac1 inhibition but could also improve Sdf1 presentation to its receptor (Matthews *et al.*, 2008; Theveneau and Mayor, 2010). Altogether, these findings and previous work on cell migration strongly suggest that small GTPases may be molecules of choice to convert a wide range of external inputs into directional information driving collective cell migration (Parsons *et al.*, 2010; Ridley *et al.*, 2003; Theveneau and Mayor, 2010).

Recent advances make it possible to integrate Contact-Inhibition of Locomotion, EMT, positive, and negative taxis into a global model of CNC behavior during migration (see Fig. 4). However, information on the downstream effectors mediating all these signaling pathways and their putative interconnections are still sketchy. In addition, the proportion of cells able to exhibit Contact-Inhibition of Locomotion or undergo efficient chemotaxis remains unknown. Moreover, recent work on trunk NC cells in the chicken embryo showed that CXCR4-positive and CXCR4-negative subpopulations are driven to different locations where they give rise to different cell types (Kasemeier-Kulesa *et al.*, 2010). These data indicate that the migratory NC population is likely to be composed of several subpopulations of specific abilities. It also reinforces the idea proposed for trunk melanocytes that ability to invade a specific region and fate restriction may somehow be linked (Erickson and Goins, 1995). Despite being supported by observations made on trunk NC cells, these questions have yet to be addressed at the cephalic level.

Further investigation is needed to assess the level of diversity of the CNC population and to fully understand how cells integrate information at the molecular level to make choices resulting in their targeting to precise locations.

ACKNOWLEDGMENTS

We thank Rachel Moore, Mae Woods, and Manuela Melchionda for comments on the manuscript.

LITERATURE CITED

- Abercrombie M, Heaysman JE. 1953. Observations on the social behaviour of cells in tissue culture. I. Speed of movement of chick heart fibroblasts in relation to their mutual contacts. *Exp Cell Res* 5:111-131.
- Adams RH, Diella F, Hennig S, Helmbacher F, Deutsch U, Klein R. 2001. The cytoplasmic domain of the ligand ephrinB2 is required for vascular morphogenesis but not cranial neural crest migration. *Cell* 104:57-69.
- Alfandari D, Cousin H, Gaultier A, Smith K, White JM, Darribere T, DeSimone DW. 2001. Xenopus ADAM 13 is a metalloprotease required for cranial neural crest-cell migration. *Curr Biol* 11:918-930.
- Alfandari D, Wolfsberg TG, White JM, DeSimone DW. 1997. ADAM 13: A novel ADAM expressed in somitic mesoderm and neural crest cells during *Xenopus laevis* development. *Dev Biol* 182:314-330.
- Barlow AJ, Wallace AS, Thapar N, Burns AJ. 2008. Critical numbers of neural crest cells are required in the pathways from the neural tube to the foregut to ensure complete enteric nervous system formation. *Development* 135:1681-1691.
- Belmadani A, Jung H, Ren D, Miller RJ. 2009. The chemokine SDF-1/CXCL12 regulates the migration of melanocyte progenitors in mouse hair follicles. *Differentiation* 77:395-411.
- Belmadani A, Tran PB, Ren D, Assimacopoulos S, Grove EA, Miller RJ. 2005. The chemokine stromal cell-derived factor-1 regulates the migration of sensory neuron progenitors. *J Neurosci* 25:3995-4003.
- Burger JA, Kipps TJ. 2006. CXCR4: A key receptor in the crosstalk between tumor cells and their micro-environment. *Blood* 107:1761-1767.
- Busillo JM, Benovic JL. 2007. Regulation of CXCR4 signaling. *Biochim Biophys Acta* 1768:952-963.
- Cai DH, Brauer PR. 2002. Synthetic matrix metalloproteinase inhibitor decreases early cardiac neural crest migration in chicken embryos. *Dev Dyn* 224:441-449.
- Cai DH, Vollberg TM Sr, Hahn-Dantona E, Quigley JP, Brauer PR. 2000. MMP-2 expression during early avian cardiac and neural crest morphogenesis. *Anat Rec* 259:168-179.
- Cai H, Kratzschmar J, Alfandari D, Hunnicutt G, Blobel CP. 1998. Neural crest-specific and general expression of distinct metalloprotease-disintegrins in early *Xenopus laevis* development. *Dev Biol* 204:508-524.
- Cantemir V, Cai DH, Reedy MV, Brauer PR. 2004. Tissue inhibitor of metalloproteinase-2 (TIMP-2) expression during cardiac neural crest cell migration and its role in proMMP-2 activation. *Dev Dyn* 231:709-719.
- Carey DJ. 1997. Syndecans: Multifunctional cell-surface co-receptors. *Biochem J* 327(Pt 1):1-16.
- Carmona-Fontaine C, Matthews HK, Kuriyama S, Moreno M, Dunn GA, Parsons M, Stern CD, Mayor R. 2008. Contact inhibition of locomotion in vivo controls neural crest directional migration. *Nature* 456:957-961.
- Charnaux N, Brule S, Hamon M, Chaigneau T, Saffar L, Prost C, Lievre N, Gattegno L. 2005. Syndecan-4 is a signaling molecule for stromal cell-derived factor-1 (SDF-1)/CXCL12. *FEBS J* 272:1937-1951.
- Cohen MM Jr. 1989. Syndromology: An updated conceptual overview. I-VI. *Int J Oral Maxillofac Surg* 18:216-222.

- Cohen MM Jr. 1990. Syndromology: An updated conceptual overview. VII-X. *Int J Oral Maxillofac Surg* 19:26-32.
- Couly GF, Coltey PM, Le Douarin NM. 1993. The triple origin of skull in higher vertebrates: A study in quail-chick chimeras. *Development* 117:409-429.
- Covington MD, Bayless KJ, Burghardt RC, Davis GE, Parrish AR. 2005. Ischemia-induced cleavage of cadherins in NRK cells: Evidence for a role of metalloproteinases. *Am J Physiol Renal Physiol* 289:F280-F288.
- Covington MD, Burghardt RC, Parrish AR. 2006. Ischemia-induced cleavage of cadherins in NRK cells requires MT1-MMP (MMP-14). *Am J Physiol Renal Physiol* 290:F43-F51.
- Creuzet S, Vincent C, Couly G. 2005. Neural crest derivatives in ocular and periocular structures. *Int J Dev Biol* 49:161-171.
- D'Amico-Martel A, Noden DM. 1983. Contributions of placodal and neural crest cells to avian cranial peripheral ganglia. *Am J Anat* 166:445-468.
- David NB, Sapede D, Saint-Etienne L, Thisse C, Thisse B, Dambly-Chaudiere C, Rosa FM, Ghysen A. 2002. Molecular basis of cell migration in the fish lateral line: Role of the chemokine receptor CXCR4 and of its ligand SDF1. *Proc Natl Acad Sci USA* 99:16297-16302.
- Davy A, Aubin J, Soriano P. 2004. Ephrin-B1 forward and reverse signaling are required during mouse development. *Genes Dev* 18:572-583.
- De Bellard ME, Rao Y, Bronner-Fraser M. 2003. Dual function of Slit2 in repulsion and enhanced migration of trunk, but not vagal, neural crest cells. *J Cell Biol* 162:269-279.
- del Barrio MG, Nieto MA. 2002. Overexpression of Snail family members highlights their ability to promote chick neural crest formation. *Development* 129:1583-1593.
- Dickson BJ, Gilestro GF. 2006. Regulation of commissural axon pathfinding by slit and its Robo receptors. *Annu Rev Cell Dev Biol* 22:651-675.
- Druckenbrod NR, Epstein ML. 2007. Behavior of enteric neural crest-derived cells varies with respect to the migratory wavefront. *Dev Dyn* 236:84-92.
- Duong TD, Erickson CA. 2004. MMP-2 plays an essential role in producing epithelial-mesenchymal transformations in the avian embryo. *Dev Dyn* 229:42-53.
- Edwards DR, Handsley MM, Pennington CJ. 2008. The ADAM metalloproteinases. *Mol Aspects Med* 29:258-289.
- Eickholt BJ, Mackenzie SL, Graham A, Walsh FS, Doherty P. 1999. Evidence for collapsin-1 functioning in the control of neural crest migration in both trunk and hindbrain regions. *Development* 126:2181-2189.
- Erickson CA. 1985. Control of neural crest cell dispersion in the trunk of the avian embryo. *Dev Biol* 111:138-157.
- Erickson CA. 1988. Control of pathfinding by the avian trunk neural crest. *Development* 103(Suppl):63-80.
- Erickson CA, Goins TL. 1995. Avian neural crest cells can migrate in the dorsolateral path only if they are specified as melanocytes. *Development* 121:915-924.
- Erickson CA, Olivier KR. 1983. Negative chemotaxis does not control quail neural crest cell dispersion. *Dev Biol* 96:542-551.
- Evans DJ, Noden DM. 2006. Spatial relations between avian craniofacial neural crest and paraxial mesoderm cells. *Dev Dyn* 235:1310-1325.
- Gammill LS, Gonzalez C, Bronner-Fraser M. 2007. Neuropilin 2/semaphorin 3F signaling is essential for cranial neural crest migration and trigeminal ganglion condensation. *Dev Neurobiol* 67:47-56.
- Ganju RK, Brubaker SA, Meyer J, Dutt P, Yang Y, Qin S, Newman W, Groopman JE. 1998. The α -chemokine, stromal cell-derived factor-1 α , binds to the transmembrane G-protein-coupled CXCR-4 receptor and activates multiple signal transduction pathways. *J Biol Chem* 273:23169-23175.
- Gaultier A, Cousin H, Darribere T, Alfandari D. 2002. ADAM13 disintegrin and cysteine-rich domains bind to the second heparin-binding domain of fibronectin. *J Biol Chem* 277:23336-23344.
- Giambernardi TA, Sakaguchi AY, Gluhak J, Pavlin D, Troyer DA, Das G, Rodeck U, Klebe RJ. 2001. Neutrophil collagenase (MMP-8) is expressed during early development in neural crest cells as well as in adult melanoma cells. *Matrix Biol* 20:577-587.
- Giannelli G, Falk-Marzillier J, Schiraldi O, Stetler-Stevenson WG, Quaranta V. 1997. Induction of cell migration by matrix metalloproteinase-2 cleavage of laminin-5. *Science* 277:225-228.
- Graveson AC, Smith MM, Hall BK. 1997. Neural crest potential for tooth development in a urodele amphibian: Developmental and evolutionary significance. *Dev Biol* 188:34-42.
- Gross JB, Hanken J. 2005. Cranial neural crest contributes to the bony skull vault in adult *Xenopus laevis*: Insights from cell labeling studies. *J Exp Zool B Mol Dev Evol* 304:169-176.
- Gross JB, Hanken J. 2008. Review of fate-mapping studies of osteogenic cranial neural crest in vertebrates. *Dev Biol* 317:389-400.
- Hall B. 2008. The neural crest and neural crest cells in vertebrate development and evolution, 2nd ed. New York: Springer.
- Hanken J, Gross JB. 2005. Evolution of cranial development and the role of neural crest: Insights from amphibians. *J Anat* 207:437-446.
- Harrison M, Abu-Elmagd M, Grocott T, Yates C, Gavrilovic J, Wheeler GN. 2004. Matrix metalloproteinase genes in *Xenopus* development. *Dev Dyn* 231:214-220.

- Hartland SN, Murphy F, Aucott RL, Abergel A, Zhou X, Waung J, Patel N, Bradshaw C, Collins J, Mann D, Benyon RC, Iredale JP. 2009. Active matrix metalloproteinase-2 promotes apoptosis of hepatic stellate cells via the cleavage of cellular N-cadherin. *Liver Int* 29:966-978.
- Helms JA, Cordero D, Tapadia MD. 2005. New insights into craniofacial morphogenesis. *Development* 132:851-861.
- Helms JA, Schneider RA. 2003. Cranial skeletal biology. *Nature* 423:326-331.
- Ho L, Symes K, Yordan C, Gudas LJ, Mercola M. 1994. Localization of PDGF A and PDGFR alpha mRNA in *Xenopus* embryos suggests signalling from neural ectoderm and pharyngeal endoderm to neural crest cells. *Mech Dev* 48:165-174.
- Imai S, Okuno M, Moriwaki H, Muto Y, Murakami K, Shudo K, Suzuki Y, Kojima S. 1997. 9,13-di-*cis*-Retinoic acid induces the production of tPA and activation of latent TGF- β via RAR alpha in a human liver stellate cell line. *LI90. FEBS Lett* 411:102-106.
- Jia L, Cheng L, Raper J. 2005. Slit/Robo signaling is necessary to confine early neural crest cells to the ventral migratory pathway in the trunk. *Dev Biol* 282:411-421.
- Jiang Y, Liu MT, Gershon MD. 2003. Netrins and DCC in the guidance of migrating neural crest-derived cells in the developing bowel and pancreas. *Dev Biol* 258:364-384.
- Johnston MC, Noden DM, Hazelton RD, Coulombre JL, Coulombre AJ. 1979. Origins of avian ocular and periocular tissues. *Exp Eye Res* 29:27-43.
- Jones MC. 1990. The neurocristopathies: Reinterpretation based upon the mechanism of abnormal morphogenesis. *Cleft Palate J* 27:136-140.
- Kasemeier-Kulesa JC, McLennan R, Romine MH, Kulesa PM, Lefcort F. 2010. CXCR4 controls ventral migration of sympathetic precursor cells. *J Neurosci* 30:13078-13088.
- Koestner U, Shnitsar I, Linnemannstons K, Hufton AL, Borchers A. 2008. Semaphorin and neuropilin expression during early morphogenesis of *Xenopus laevis*. *Dev Dyn* 237:3853-3863.
- Kulesa PM, Bailey CM, Kasemeier-Kulesa JC, McLennan R. 2010. Cranial neural crest migration: New rules for an old road. *Dev Biol* 344:543-554.
- Kulesa PM, Fraser SE. 2000. In ovo time-lapse analysis of chick hindbrain neural crest cell migration shows cell interactions during migration to the branchial arches. *Development* 127:1161-1172.
- Kulesa PM, Teddy JM, Stark DA, Smith SE, McLennan R. 2008. Neural crest invasion is a spatially-ordered progression into the head with higher cell proliferation at the migratory front as revealed by the photoactivatable protein KikGR. *Dev Biol* 316:275-287.
- Kuriyama S, Mayor R. 2008. Molecular analysis of neural crest migration. *Philos Trans R Soc Lond B Biol Sci* 363:1349-1362.
- Le Douarin N, Kalcheim C. 1999. The neural crest, 2nd ed. Cambridge, UK: Cambridge University Press. xxiii, 445 p.
- Le Lievre CS. 1978. Participation of neural crest-derived cells in the genesis of the skull in birds. *J Embryol Exp Morphol* 47:17-37.
- Lee VM, Sechrist JW, Luetolf S, Bronner-Fraser M. 2003. Both neural crest and placode contribute to the ciliary ganglion and oculomotor nerve. *Dev Biol* 263:176-190.
- Lumsden A, Guthrie S. 1991. Alternating patterns of cell surface properties and neural crest cell migration during segmentation of the chick hindbrain. *Dev Suppl* 2:9-15.
- Lumsden A, Sprawson N, Graham A. 1991. Segmental origin and migration of neural crest cells in the hindbrain region of the chick embryo. *Development* 113:1281-1291.
- Lumsden AG. 1988. Spatial organization of the epithelium and the role of neural crest cells in the initiation of the mammalian tooth germ. *Development* 103(Suppl):155-169.
- Lumsden AG, Buchanan JA. 1986. An experimental study of timing and topography of early tooth development in the mouse embryo with an analysis of the role of innervation. *Arch Oral Biol* 31:301-311.
- Martino MM, Hubbell JA. 2010. The 12th-14th type III repeats of fibronectin function as a highly promiscuous growth factor-binding domain. *FASEB J* 24:4711-4721.
- Matthews HK, Marchant L, Carmona-Fontaine C, Kuriyama S, Larrain J, Holt MR, Parsons M, Mayor R. 2008. Directional migration of neural crest cells in vivo is regulated by Syndecan-4/Rac1 and non-canonical Wnt signaling/RhoA. *Development* 135:1771-1780.
- Mayor R, Carmona-Fontaine C. 2010. Keeping in touch with contact inhibition of locomotion. *Trends Cell Biol* 20:319-328.
- McCusker C, Cousin H, Neuner R, Alfandari D. 2009. Extracellular cleavage of cadherin-11 by ADAM metalloproteases is essential for *Xenopus* cranial neural crest cell migration. *Mol Biol Cell* 20:78-89.
- McLennan R, Kulesa PM. 2007. In vivo analysis reveals a critical role for neuropilin-1 in cranial neural crest cell migration in chick. *Dev Biol* 301:227-239.
- McLennan R, Kulesa PM. 2010. Neuropilin-1 interacts with the second branchial arch microenvironment to mediate chick neural crest cell dynamics. *Dev Dyn* 239:1664-1673.
- McLennan R, Teddy JM, Kasemeier-Kulesa JC, Romine MH, Kulesa PM. 2010. Vascular endothelial growth

- factor (VEGF) regulates cranial neural crest migration in vivo. *Dev Biol* 339:114–125.
- McQuibban GA, Butler GS, Gong JH, Bendall L, Power C, Clark-Lewis I, Overall CM. 2001. Matrix metalloproteinase activity inactivates the CXC chemokine stromal cell-derived factor-1. *J Biol Chem* 276: 43503–43508.
- Mellott DO, Burke RD. 2008. Divergent roles for Eph and ephrin in avian cranial neural crest. *BMC Dev Biol* 8:56.
- Minoux M, Rijli FM. 2010. Molecular mechanisms of cranial neural crest cell migration and patterning in craniofacial development. *Development* 137:2605–2621.
- Molyneaux KA, Zinszner H, Kunwar PS, Schaible K, Stebler J, Sunshine MJ, O'Brien W, Raz E, Littman D, Wylie C, Lehmann R. 2003. The chemokine SDF1/CXCL12 and its receptor CXCR4 regulate mouse germ cell migration and survival. *Development* 130:4279–4286.
- Morriss-Kay GM. 2001. Derivation of the mammalian skull vault. *J Anat* 199:143–151.
- Neuner R, Cousin H, McCusker C, Coyne M, Alfandari D. 2009. Xenopus ADAM19 is involved in neural, neural crest and muscle development. *Mech Dev* 126:240–255.
- Noden DM, Trainor PA. 2005. Relations and interactions between cranial mesoderm and neural crest populations. *J Anat* 207:575–601.
- Olesnicki Killian EC, Birkholz DA, Artinger KB. 2009. A role for chemokine signaling in neural crest cell migration and craniofacial development. *Dev Biol* 333:161–172.
- Orr-Urtreger A, Bedford MT, Do MS, Eisenbach L, Lonai P. 1992. Developmental expression of the alpha receptor for platelet-derived growth factor, which is deleted in the embryonic lethal Patch mutation. *Development* 115:289–303.
- Orr-Urtreger A, Lonai P. 1992. Platelet-derived growth factor-A and its receptor are expressed in separate, but adjacent cell layers of the mouse embryo. *Development* 115:1045–1058.
- Osborne NJ, Begbie J, Chilton JK, Schmidt H, Eickholt BJ. 2005. Semaphorin/neuropilin signaling influences the positioning of migratory neural crest cells within the hindbrain region of the chick. *Dev Dyn* 232:939–949.
- Parsons JT, Horwitz AR, Schwartz MA. 2010. Cell adhesion: Integrating cytoskeletal dynamics and cellular tension. *Nat Rev Mol Cell Biol* 11:633–643.
- Pelletier AJ, van der Laan LJ, Hildbrand P, Siani MA, Thompson DA, Dawson PE, Torbett BE, Salomon DR. 2000. Presentation of chemokine SDF-1 alpha by fibronectin mediates directed migration of T cells. *Blood* 96:2682–2690.
- Perez-Alcala S, Nieto MA, Barbas JA. 2004. LSox5 regulates RhoB expression in the neural tube and promotes generation of the neural crest. *Development* 131:4455–4465.
- Petitclerc E, Stromblad S, von Schalscha TL, Mitjans F, Piulats J, Montgomery AM, Cheresch DA, Brooks PC. 1999. Integrin $\alpha_v\beta_3$ promotes M21 melanoma growth in human skin by regulating tumor cell survival. *Cancer Res* 59:2724–2730.
- Piloto S, Schilling TF. 2010. Ovo1 links Wnt signaling with N-cadherin localization during neural crest migration. *Development* 137:1981–1990.
- Raz E, Mahabaleshwar H. 2009. Chemokine signaling in embryonic cell migration: A fisheye view. *Development* 136:1223–1229.
- Rehimi R, Khalida N, Yusuf F, Dai F, Morosan-Puopolo G, Brand-Saberi B. 2008. Stromal-derived factor-1 (SDF-1) expression during early chick development. *Int J Dev Biol* 52:87–92.
- Ridley AJ, Schwartz MA, Burridge K, Firtel RA, Ginsberg MH, Borisy G, Parsons JT, Horwitz AR. 2003. Cell migration: Integrating signals from front to back. *Science* 302:1704–1709.
- Rodriguez D, Morrison CJ, Overall CM. 2010. Matrix metalloproteinases: What do they not do? New substrates and biological roles identified by murine models and proteomics. *Biochim Biophys Acta* 1803:39–54.
- Santiago A, Erickson CA. 2002. Ephrin-B ligands play a dual role in the control of neural crest cell migration. *Development* 129:3621–3632.
- Schatteman GC, Morrison-Graham K, van Koppen A, Weston JA, Bowen-Pope DE. 1992. Regulation and role of PDGF receptor α -subunit expression during embryogenesis. *Development* 115:123–131.
- Schoenwolf GC, Bleyl SB, Brauer PR, Francis-West PH, Larsen WJ. 2009. Larsen's human embryology, 4th ed. Philadelphia: Churchill Livingstone/Elsevier. xix,687p.
- Schwarz Q, Vieira JM, Howard B, Eickholt BJ, Ruhrberg C. 2008. Neuropilin 1 and 2 control cranial gangliogenesis and axon guidance through neural crest cells. *Development* 135:1605–1613.
- Sechrist J, Scherson T, Bronner-Fraser M. 1994. Rhombomere rotation reveals that multiple mechanisms contribute to the segmental pattern of hindbrain neural crest migration. *Development* 120: 1777–1790.
- Sechrist J, Serbedzija GN, Scherson T, Fraser SE, Bronner-Fraser M. 1993. Segmental migration of the hindbrain neural crest does not arise from its segmental generation. *Development* 118:691–703.
- Shiau CE, Bronner-Fraser M. 2009. N-cadherin acts in concert with Slit1-Robo2 signaling in regulating aggregation of placode-derived cranial sensory neurons. *Development* 136:4155–4164.
- Shoval I, Ludwig A, Kalcheim C. 2007. Antagonistic roles of full-length N-cadherin and its soluble BMP cleavage product in neural crest delamination. *Development* 134:491–501.

- Simpson MJ, Zhang DC, Mariani M, Landman KA, Newgreen DE. 2007. Cell proliferation drives neural crest cell invasion of the intestine. *Dev Biol* 302:553-568.
- Smith A, Robinson V, Patel K, Wilkinson DG. 1997. The EphA4 and EphB1 receptor tyrosine kinases and ephrin-B2 ligand regulate targeted migration of branchial neural crest cells. *Curr Biol* 7:561-570.
- Smith CL, Tallquist MD. 2010. PDGF function in diverse neural crest cell populations. *Cell Adhesion & Migration* 4:4, 561-566.
- Smith MM, Hall BK. 1990. Development and evolutionary origins of vertebrate skeletogenic and odontogenic tissues. *Biol Rev Camb Philos Soc* 65:277-373.
- Stebler J, Spieler D, Slanchev K, Molyneaux KA, Richter U, Cojocaru V, Tarabykin V, Wylie C, Kessel M, Raz E. 2004. Primordial germ cell migration in the chick and mouse embryo: The role of the chemokine SDF-1/CXCL12. *Dev Biol* 272:351-361.
- Sternlicht MD, Werb Z. 2001. How matrix metalloproteinases regulate cell behavior. *Annu Rev Cell Dev Biol* 17:463-516.
- Takakura N, Yoshida H, Ogura Y, Kataoka H, Nishikawa S. 1997. PDGFR alpha expression during mouse embryogenesis: Immunolocalization analyzed by whole-mount immunohistostaining using the monoclonal anti-mouse PDGFR alpha antibody APA5. *J Histochem Cytochem* 45:883-893.
- Taneyhill LA, Coles EG, Bronner-Fraser M. 2007. Snail2 directly represses cadherin6B during epithelial-to-mesenchymal transitions of the neural crest. *Development* 134:1481-1490.
- Teddy JM, Kulesa PM. 2004. In vivo evidence for short- and long-range cell communication in cranial neural crest cells. *Development* 131:6141-6151.
- Theveneau E, Duband JL, Altabef M. 2007. Ets-1 confers cranial features on neural crest delamination. *PLoS One* 2:e1142.
- Theveneau E, Marchant L, Kuriyama S, Gull M, Moepps B, Parsons M, Mayor R. 2010. Collective chemotaxis requires contact-dependent cell polarity. *Dev Cell* 19:39-53.
- Theveneau E, Mayor R. 2010. Integrating chemotaxis and contact-inhibition during collective cell migration: Small GTPases at work. *Small GTPases* 1:2, 1-5.
- Tomlinson ML, Guan P, Morris RJ, Fidock MD, Rejzek M, Garcia-Morales C, Field RA, Wheeler GN. 2009. A chemical genomic approach identifies matrix metalloproteinases as playing an essential and specific role in *Xenopus* melanophore migration. *Chem Biol* 16:93-104.
- Trainor PA. 2005. Specification and patterning of neural crest cells during craniofacial development. *Brain Behav Evol* 66:266-280.
- Trainor PA. 2010. Craniofacial birth defects: The role of neural crest cells in the etiology and pathogenesis of Treacher Collins syndrome and the potential for prevention. *Am J Med Genet A* 152:2984-2994.
- Trainor PA, Tam PP. 1995. Cranial paraxial mesoderm and neural crest cells of the mouse embryo: Co-distribution in the craniofacial mesenchyme but distinct segregation in branchial arches. *Development* 121:2569-2582.
- Trokovic N, Trokovic R, Partanen J. 2005. Fibroblast growth factor signalling and regional specification of the pharyngeal ectoderm. *Int J Dev Biol* 49:797-805.
- Walker MB, Trainor PA. 2006. Craniofacial malformations: Intrinsic vs extrinsic neural crest cell defects in Treacher Collins and 22q11 deletion syndromes. *Clin Genet* 69:471-479.
- Wurdak H, Ittner LM, Sommer L. 2006. DiGeorge syndrome and pharyngeal apparatus development. *Bioessays* 28:1078-1086.
- Ypsilanti AR, Zagar Y, Chedotal A. 2010. Moving away from the midline: New developments for Slit and Robo. *Development* 137:1939-1952.
- Yu HH, Moens CB. 2005. Semaphorin signaling guides cranial neural crest cell migration in zebrafish. *Dev Biol* 280:373-385.
- Yusuf F, Rehim R, Dai F, Brand-Saberi B. 2005. Expression of chemokine receptor CXCR4 during chick embryo development. *Anat Embryol (Berl)* 210:35-41.